



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

CORRESPONDENCE.

Editor American Naturalist:

In the July number of the *American Naturalist*, among his "Notes on the Cœlenterate Fauna of Woods Hole," Dr. C. W. Hargitt has defined what he considers to be a new species of tubularian hydroid, under the name of *Tubularia parasitica*. During the past summer I have been enabled to observe the development of the western representative of the genus *Corymorpha*, *C. palma*, a few facts concerning which will show, I think, that *T. parasitica* is but a young form of the *pendula* on which it was growing. It has seemed best that attention should be called to this at once to prevent future inconvenience to taxonomists.

As with *C. pendula*, the medusæ of *C. palma* are permanently attached. The eggs arise on the manubrium, break through the ectoderm when ripe, and, in quiet water, settle at once to the bottom. The egg case is adhesive, fastening to the first object with which it comes in contact. Many eggs are dropped at the same time, and often cling to the rootlets of the parent hydroid, where *T. parasitica* was found. There is no free larval stage. The embryo is able, however, to change its location very slowly, leaving a trail of perisarc behind it. Such a movement accounts for the clusters of six or eight individuals which are commonly found adhering to each other near their bases. The stem has a single central canal at first, as in *Tubularia*. The peduncles which support the medusæ appear very early. When there are not more than eight proximal tentacles, buds arise on the base of the hydroid and develop into rootlets. They are the structures, I suspect, which Dr. Hargitt has taken for "absorbent organs."

The young *Corymorpha* further agrees with *T. parasitica* in size, number of tentacles, and general aspect.

I shall describe the development of *C. palma* more fully in another connection.

HARRY BEAL TORREY.

ZOOLOGICAL LABORATORY, UNIVERSITY OF CALIFORNIA,
BERKELEY, CAL., Sept. 5, 1902.